



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CALCULUS.

133. Proposed by NELSON L. RORAY, South Jersey Institute, Bridgeton, N. J.

Integrate $\int \frac{\sqrt{1+y}}{1+y^2} dy$.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Let $1+y=z^2$.

$$\begin{aligned} \therefore \int \frac{\sqrt{1+y}}{1+y^2} dy &= \int \frac{2z^2 dz}{1+[z^2-1]^2} = [1+\sqrt{-1}] \int \frac{dz}{1-\sqrt{-1}-1+z^2\sqrt{-1}} \\ &+ [1-\sqrt{-1}] \int \frac{dz}{1+\sqrt{-1}-1-z^2\sqrt{-1}} = -[1-\sqrt{-1}] \int \frac{dz}{1+\sqrt{-1}-1-z^2} \\ &- [1+\sqrt{-1}] \int \frac{dz}{1-\sqrt{-1}-1-z^2} = u. \end{aligned}$$

Let $1+\sqrt{-1}=a^2$, $1-\sqrt{-1}=b^2$.

$$\begin{aligned} \therefore u &= -\frac{1}{a^3} \int \left[\frac{1}{a+z} + \frac{1}{a-z} \right] dz - \frac{1}{b^3} \int \left[\frac{1}{b+z} + \frac{1}{b-z} \right] dz \\ &= \frac{1}{a^3} \log \left[\frac{a-z}{a+z} \right] + \frac{1}{b^3} \log \left[\frac{b-z}{b+z} \right] + C. \end{aligned}$$

$$\begin{aligned} \therefore u &= \left[\frac{1}{1+\sqrt{-1}} \right]^{\frac{3}{2}} \log \left[\frac{\sqrt{1+\sqrt{-1}}-1}{\sqrt{1+\sqrt{-1}}+1} \right] \\ &+ \left[\frac{1}{1-\sqrt{-1}} \right]^{\frac{3}{2}} \log \left[\frac{\sqrt{1-\sqrt{-1}}-1}{\sqrt{1-\sqrt{-1}}+1} \right] + C. \end{aligned}$$

Also solved by F. P. MATZ. An incorrect solution was received from H. C. WHITAKER.

134. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

To find the curve for which the sum of that part of the tangent, lying between the point of contact and the axis of abscissas, and the corresponding ordinate is constant= c , and which passes through the point (a, b) .

Solution by F. P. MATZ, Sc. D., Ph. D., Defiance College, Defiance, O.; G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa.; COOPER D. SCHMIDT, A. M., University of Tennessee, Knoxville, Tenn.; and the PROPOSER.

According to the conditions of the problem, *Ordinate* + *Tangent* = *Constant*.

That is, $y + y \sqrt{\left(\frac{dx}{dy}\right)^2 + 1} = C \dots [1]$.

Therefore, $dx = \frac{\sqrt{C^2 - 2Cy}}{y} dy \dots [2]$.

or, $x = 2\sqrt{C^2 - 2Cy} + C \log \left[\frac{\sqrt{C^2 - 2Cy} - C}{\sqrt{C^2 - 2Cy} + C} \right] + C' \dots [3]$.

For $x=a$ and $y=b$, as per the problem, [3] gives the required equation

$$x - a = 2\{\sqrt{C^2 - 2Cy} - \sqrt{C^2 - 2bC}\} \\ + C \log \left[\frac{\sqrt{C^2 - 2Cy} - C}{\sqrt{C^2 - 2Cy} + C} \cdot \frac{\sqrt{C^2 - 2bC} - C}{\sqrt{C^2 - 2bC} + C} \right] \dots [4].$$

Also solved by *L. C. WALKER* and *H. C. WHITAKER*. Professor Matz gave a second solution using polar co-ordinates.

135. Proposed by *COOPER D. SCHMITT*, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

To find the equation of the evolute of the common catenary

$$y = (\tfrac{1}{2}c)(e^{x/c} + e^{-x/c}).$$

Solution by *G. B. M. ZEER*, A. M., Ph. D., The Temple College, Philadelphia, Pa., and *M. E. GRABER*, Heidelberg University, Tiffin, O.

$$\frac{dy}{dx} = \tfrac{1}{2}(e^{x/c} - e^{-x/c}), \quad \frac{d^2y}{dx^2} = \frac{1}{2c}(e^{x/c} + e^{-x/c}).$$

$\therefore x - m + (y - n)dy/dx = 0$, $1 + (dy/dx)^2 + (y - n)d^2y/dx^2 = 0$ become

$$x - m + \tfrac{1}{2}c(e^{2x/c} - e^{-2x/c}) - \tfrac{1}{2}n(e^{x/c} - e^{-x/c}) = 0 \dots (1).$$

$$1 + \tfrac{1}{4}(e^{x/c} - e^{-x/c})^2 + \tfrac{1}{4}(e^{x/c} + e^{-x/c})^2 - \frac{n}{2c}(e^{x/c} + e^{-x/c}) = 0 \dots (2).$$

$$\text{From (2), } e^{4x/c} - \frac{n}{c}e^{3x/c} + 2e^{2x/c} - \frac{n}{c}e^{x/c} + 1 = 0.$$

Let $e^{x/c} = z$.

$$\therefore z^4 - \frac{n}{c}z^3 + 2z^2 - \frac{n}{c}z + 1 = 0, \quad (z^2 + 1)(z^2 - \frac{n}{c}z + 1) = 0.$$

$$\therefore z = \frac{n \pm \sqrt{(n^2 - 4c^2)}}{2c} = e^{x/c} \text{ and } \frac{n \mp \sqrt{(n^2 - 4c^2)}}{2c} = e^{-x/c}.$$

$$\therefore x = c \log \left(\frac{n \pm \sqrt{(n^2 - 4c^2)}}{2c} \right).$$